arXiv:1109.2157v1 [math.PR] 9 Sep 2011

# Spectral Conditions for Strong Local Nondeterminism and Exact Hausdorff Measure of Ranges of Gaussian Random Fields

Nana Luan
University of International Business and Economics, Beijing

Yimin Xiao [*]
Michigan State University

September 13, 2011

## Abstract

Let $X = \{X(t), t \in \mathbb{R}^N\}$ be a Gaussian random field with values in $\mathbb{R}^d$ defined by

$$X(t) = \big(X_1(t), \ldots, X_d(t)\big), \qquad t \in \mathbb{R}^N,$$

where $X_1, \ldots, X_d$ are independent copies of a real-valued, centered, anisotropic Gaussian random field $X_0$ which has stationary increments and the property of strong local nondeterminism. In this paper we determine the exact Hausdorff measure function for the range $X([0,1]^N)$.

We also provide a sufficient condition for a Gaussian random field with stationary increments to be strongly locally nondeterministic. This condition is given in terms of the spectral measures of the Gaussian random fields which may contain either an absolutely continuous or discrete part. This result strengthens and extends significantly the related theorems of Berman (1973, 1988), Pitt (1978) and Xiao (2007, 2009), and will have wider applicability beyond the scope of the present paper.

Running head: Strong Local Nondeterminism and Hausdorff Measure of Gaussian Random Fields

2010 AMS Classification numbers: 60G15, 60G17, 60G60, 28A80.

Key words: Gaussian random fields, strong local nondeterminism, spectral condition, anisotropy, Hausdorff dimension, Hausdorff measure.

## 1 Introduction

Let $X = \{X(t), t \in \mathbb{R}^N\}$ be a Gaussian random field with values in $\mathbb{R}^d$, where

$$X(t) = \big(X_1(t), \ldots, X_d(t)\big), \qquad t \in \mathbb{R}^N. \tag{1.1}$$

---

[*] Research partially supported by NSF grant DMS-1006903.

For brevity we call $X$ an $(N, d)$-Gaussian random field. Sample path properties of $X$ such as the Hausdorff dimensions of the range $X([0,1]^N) = \{X(t) : t \in [0,1]^N\}$, the graph $\mathrm{Gr}X([0,1]^N) = \{(t, X(t)) : t \in [0,1]^N\}$ and the level set $X^{-1}(x) = \{t \in \mathbb{R}^N : X(t) = x\}$ $(x \in \mathbb{R}^d)$ have been studied by many authors under various assumptions on the coordinate processes $X_1, \ldots, X_d$. We refer to Adler (1981), Kahane (1985) and Xiao (2007, 2009) for further information.

In the cases when $X_1, \ldots, X_d$ are independent copies of an approximately *isotropic* Gaussian random field $X_0$ [a typical example is fractional Brownian motion], the problems for finding the exact Hausdorff measure functions for $X([0,1]^N)$, $\mathrm{Gr}X([0,1]^N)$ and $X^{-1}(x)$ have been investigated by Talagrand (1995, 1998), Xiao (1996, 1997a, 1997b), Baraka and Mountford (2008, 2011).

The main objective of this paper is to study the exact Hausdorff measure of the range of Gaussian random fields which are anisotropic in the time-variable. More specifically, we consider an $(N, d)$-Gaussian random field $X = \{X(t), t \in \mathbb{R}^N\}$ whose coordinate processes $X_1, \ldots, X_d$ in (1.1) are independent copies of a centered, real-valued Gaussian field $X_0$ with stationary increments and $X_0(0) = 0$ almost surely; and we assume there exists a constant vector $H = (H_1, \cdots, H_N) \in (0,1)^N$ such that the following conditions hold:

(C1). There exists a positive constant $c_{1,1} \geq 1$ such that

$$c_{1,1}^{-1}\, \rho(s,t)^2 \leq \mathbb{E}\left(X_0(s) - X_0(t)\right)^2 \leq c_{1,1}\, \rho(s,t)^2 \quad \text{for all } s, t \in [0,1]^N,$$

where $\rho(s,t)$ is the metric on $\mathbb{R}^N$ defined by

$$\rho(s,t) = \sum_{j=1}^{N} |s_j - t_j|^{H_j}, \qquad \forall s, t \in \mathbb{R}^N.$$

(C2). There exists a positive constant $c_{1,2}$ such that for all integers $n \geq 1$ and all $u, t^1, \cdots, t^n \in [0,1]^N$, we have

$$\mathrm{Var}(X_0(u) | X_0(t^1), \cdots, X_0(t^n)) \geq c_{1,2} \min_{0 \leq k \leq n} \rho(u, t^k)^2, \qquad (t^0 = 0).$$

Section 2 below provides a way to construct a large class of Gaussian random fields with stationary increments that satisfy (C1) and (C2). Further examples can be found in Xiao (2009) and Luan and Xiao (2010). Under Condition (C1), the $(N, d)$-Gaussian random field $X$ has a version which has continuous sample functions on $[0,1]^N$ almost surely. Henceforth we will assume without loss of generality that the Gaussian random field $X$ has continuous sample paths. When $\{X_0(t), t \in \mathbb{R}^N\}$ satisfies (C2), we say that $X_0$ has the property of strong local nondeterminism in metric $\rho$ on $[0,1]^N$.

Xiao (2009) proved that, if Condition (C1) holds, then with probability 1,

$$\dim_{\mathrm{H}} X\big([0,1]^N\big) = \min\left\{d;\ \sum_{j=1}^{N} \frac{1}{H_j}\right\}, \tag{1.2}$$

where $\sum_{j=1}^{0} \frac{1}{H_j} := 0$. In the above, $\dim_{\mathrm{H}}$ denotes Hausdorff dimension [cf. Kahane (1985) or Falconer (1990)]. Further analytic and fractal properties of Gaussian random fields which satisfy Conditions (C1) and (C2) have been studied by Xiao (2009), Biermé *et al.* (2009), Luan and Xiao (2010), Meerschaert *et al.* (2011) [see also Benassi *et al.* (1997), Ayache and Xiao (2005), Wu and Xiao (2009, 2011) for related results].

The first objective of this paper is to refine (1.2) by determining the exact Hausdorff measure function for the range $X([0,1]^N)$.

**Theorem 1.1** *Let $X = \{X(t), t \in \mathbb{R}^N\}$ be an $(N,d)$-Gaussian random field with stationary increments defined by (1.1), where $X_1, \ldots, X_d$ are independent copies of a centered, real-valued Gaussian field $X_0$ with stationary increments and $X_0(0) = 0$. We assume that $X_0$ satisfies Conditions (C1) and (C2). If $d > \sum_{j=1}^N H_j^{-1}$, then we have*

$$0 < \varphi_1\text{-}m(X([0,1]^N)) < \infty \quad a.s.,$$

*where $\varphi_1$ is the function*

$$\varphi_1(r) = r^{\sum_{j=1}^N H_j^{-1}} \log\log \frac{1}{r}$$

*and $\varphi_1$-$m$ is the corresponding Hausdorff measure.*

The following remark is concerned with the cases not covered by Theorem 1.1.

**Remark 1.2**

- If $d < \sum_{j=1}^N H_j^{-1}$, then Theorem 8.2 in Xiao (2009) implies that $X([0,1]^N)$ a.s. has interior points and hence has positive $d$-dimensional Lebesgue measure. In this case, Wu and Xiao (2011) showed that $X$ has a jointly continuous local time and provides a lower bound for the exact Hausdorff measure (in the metric $\rho$) of the level set $X^{-1}(x)$. For fractional Brownian motion and some other isotropic Gaussian random fields, the exact Hausdorff measure function for $X^{-1}(x)$ has been determined by Xiao (1997b) and Baraka and Mountford (2011). However, no such result has been established for *anisotropic* Gaussian random fields.
- If $d = \sum_{j=1}^N H_j^{-1}$, then $\dim_{\mathrm{H}} X([0,1]^N) = d$ a.s. The problem to determine the exact Hausdorff measure function for $X([0,1]^N)$ in this "critical case" is open and is certainly a deeper question.

It will become clear that the proof of Theorem 1.1 relies crucially on Condition (C2)– the property of strong local nondeterminism, which is useful for studying many other sample path and statistical properties of Gaussian random fields [cf. Xiao (2009), Xue and Xiao (2011)]. The second objective of this paper is to provide a rather general condition for a Gaussian random field with stationary increments to satisfy both Conditions (C1) and (C2). This condition is given in terms of the spectral measures of the Gaussian random fields which may contain either an absolutely continuous or a discrete part. Theorem 2.4 extends the related theorems of Berman (1973, 1988), Pitt (1978) and Xiao (2007, 2009), which will have wider applicability beyond the scope of the present paper. For example, we can apply this theorem to prove that the solution of a fractional stochastic heat equation on the circle $\mathbb{S}_1$ [see

Tindel, Tudor and Viens (2004), Nualart and Viens (2009)] has the property of strong local nondeterminism in the space variable (at fixed time $t$). Hence fine properties of the sample functions of the solution can be obtained by using the results in Monrad and Rootzén (1995), Xiao (2009), Luan and Xiao (2010), and Meerschaert, Wang and Xiao (2011). Similarly, we can show that the spherical fractional Brownian motion on $\mathbb{S}_1$ introduced by Istas (2005) is also strongly locally nondeterministic. Both of these processes share local properties with ordinary fractional Brownian motion with appropriate Hurst indices. Details of these results will be given elsewhere.

The rest of this paper is organized as follows. Section 2 gives a sufficient condition for a Gaussian random field with stationary increments to be strongly locally nondeterministic. Section 3 is concerned with the exact Hausdorff measure function for the range of $X$. After recalling the definition of Hausdorff measure and its basic properties, and establishing some estimates, we prove Theorem 1.1.

We end the Introduction with some notation. The inner product of $s,\, t \in \mathbb{R}^N$ is denoted by $\langle s, t\rangle$ and the Euclidean norm of $t \in \mathbb{R}^N$ is denoted $\|t\|$. Given two points $s = (s_1, \ldots, s_N) \in \mathbb{R}^N$ and $t = (t_1, \ldots, t_N) \in \mathbb{R}^N$, $s \le t$ (resp. $s < t$) means that $s_i \le t_i$ (resp. $s_i < t_i$) for all $1 \le i \le N$. When $s \le t$, we use $[s, t]$ to denote the $N$-dimensional interval (or rectangle) $[s, t] = \prod_{i=1}^{N}[s_i, t_i]$. For any $T \subseteq \mathbb{R}^N$, $f(s) \asymp g(s)$ means the ratio $f(s)/g(s)$ is bounded from below and above by positive and finite constants which are independent of $s \in T$.

Throughout this paper we will use $c$ to denote an unspecified positive and finite constant which may not be the same in each occurrence. More specific constants in Section $i$ are numbered as $c_{i,1}, c_{i,2}, \ldots$.

**Acknowledgement** This paper was written while Nana Luan was visiting Department of Statistics and Probability, Michigan State University (MSU) with the support of a grant from China Scholarship Council (CSC). She thanks MSU for the good working condition and CSC for the financial support.

The authors thank the referees for their carefully reading of the manuscript and their helpful comments.

## 2 Spectral condition for strong local nondeterminism of Gaussian fields with stationary increments

One of the major difficulties in studying the probabilistic, analytic or statistical properties of Gaussian random fields is the complexity of their dependence structures. In many circumstances, the properties of local nondeterminism can help us to overcome this difficulty so that many elegant and deep results for Brownian motion can be extended to Gaussian random fields; see Berman (1973, 1988), Pitt (1978) and Xiao (2007, 2009) for further information. Hence, for a given Gaussian random field, it is an interesting question to determine whether it satisfies certain forms of local nondeterminism. In this section we provide a general sufficient condition for a Gaussian random field with stationary increments to satisfy Conditions (C1) and (C2).

Let $X_0 = \{X_0(t), t \in \mathbb{R}^N\}$ be a real-valued, centered Gaussian random field with stationary increments and $X_0(0) = 0$. We assume that $X_0$ has continuous covariance function $R(s, t) =$

$\mathbb{E}[X(s)X(t)]$. According to Yaglom (1957), $R(s,t)$ can be represented as

$$R(s,t) = \int_{\mathbb{R}^N} \big(e^{i\langle s,\lambda\rangle} - 1\big)\big(e^{-i\langle t,\lambda\rangle} - 1\big)\, F(d\lambda) + \langle s, Mt\rangle\,, \tag{2.1}$$

where $M$ is an $N \times N$ non-negative definite matrix and $F(d\lambda)$ is a nonnegative symmetric measure on $\mathbb{R}^N\backslash\{0\}$ satisfying

$$\int_{\mathbb{R}^N} \frac{\|\lambda\|^2}{1+\|\lambda\|^2}\, F(d\lambda) < \infty. \tag{2.2}$$

In analogy to the stationary case, the measure $F$ is called the spectral measure of $X_0$. If $F$ is absolutely continuous with respect to the Lebesgue measure in $\mathbb{R}^N$, its density $f$ will be called the spectral density of $X_0$.

It follows from (2.1) that $X_0$ has the following stochastic integral representation:

$$X_0(t) \stackrel{d}{=} \int_{\mathbb{R}^N} \big(e^{i\langle t,\lambda\rangle} - 1\big)\, W(d\lambda) + \langle Y, t\rangle\,, \tag{2.3}$$

where $\stackrel{d}{=}$ means equality of all finite dimensional distributions, $Y$ is an $N$-dimensional Gaussian random vector with mean 0 and covariance matrix $M$, $W(d\lambda)$ is a centered complex-valued Gaussian random measure which is independent of $Y$ and satisfies

$$\mathbb{E}\big(W(A)\overline{W(B)}\big) = F(A\cap B) \quad \text{and} \quad W(-A) = \overline{W(A)}$$

for all Borel sets $A, B \subseteq \mathbb{R}^N$ with finite $F$-measure. The above properties of $W(d\lambda)$ ensures that the stochastic integral in (2.3) is real-valued. The spectral measure $F$ is called the control measure of $W$. Since the linear term $\langle Y, t\rangle$ in (2.3) will not have any effect on the problems considered in this paper, we will from now on assume $Y = 0$. This is equivalent to assuming $M = 0$ in (2.1). Consequently, for any $h \in \mathbb{R}^N$ we have

$$\sigma^2(h) \triangleq \mathbb{E}\big(X_0(t+h) - X_0(t)\big)^2 = 2\int_{\mathbb{R}^N} \big(1 - \cos\langle h, \lambda\rangle\,\big)\, F(d\lambda). \tag{2.4}$$

It is important to note that $\sigma^2(h)$ is a negative definite function in the sense of I. J. Schoenberg, which is determined by the spectral measure $F$. See Berg and Forst (1975) for more information on negative definite functions. If the function $\sigma^2(h)$ depends only on $\|h\|$, then $X_0$ is called an isotropic random field. More generally, if $\sigma^2(h) \asymp \phi(\|h\|)$ in a neighborhood of $h = 0$ for some nonnegative function $\phi$, then $X_0$ is called approximately isotropic.

Various centered Gaussian random fields with stationary increments can be constructed by choosing appropriate spectral measures $F$. For the well known fractional Brownian motion $B^H = \{B^H(t), t \in \mathbb{R}^N\}$ of Hurst index $H \in (0,1)$, its spectral measure has a density function

$$f_H(\lambda) = c(H,N)\frac{1}{\|\lambda\|^{2H+N}}, \tag{2.5}$$

where $c(H,N) > 0$ is a normalizing constant such that $\sigma^2(h) = \|h\|^{2H}$. Since $\sigma^2(h)$ depends on $\|h\|$ only, the increments of $B^H$ are isotropic and stationary. Examples of approximately isotropic Gaussian fields with stationary increments can be found in Xiao (2007).

A typical example of anisotropic Gaussian random field with stationary increments can be constructed by choosing the spectral density

$$f(\lambda) = \frac{1}{\left(\sum_{j=1}^N |\lambda_j|^{H_j}\right)^{2+Q}}, \qquad \forall \lambda \in \mathbb{R}^N \backslash \{0\}, \tag{2.6}$$

where the constants $H_j \in (0, 1)$ for $j = 1, \ldots, N$ and $Q = \sum_{j=1}^N H_j^{-1}$. This notation will be fixed throughout the rest of the paper.

It can be verified that $f(\lambda)$ in (2.6) satisfies (2.2) and the corresponding Gaussian random field $X_0$ has stationary increments. In the special case when $H_1 = \cdots = H_N = H$, (2.6) is very similar to (2.5). Consequently, $X_0$ shares many properties with fractional Brownian motion.

In general, $X_0$ with spectral density (2.6) is anisotropic in the sense that the sample function $X_0(t)$ has different geometric and probabilistic characteristics along different directions. This gives more flexibility from modeling point of view. Moreover, $X_0$ is operator-self-similar with exponent $A = (a_{ij})$, where $a_{ii} = H_i^{-1}$ and $a_{ij} = 0$ if $i \neq j$. The latter means that for any constant $c > 0$,

$$\left\{X_0(c^A t),\, t \in \mathbb{R}^N\right\} \overset{d}{=} \left\{c\, X_0(t),\, t \in \mathbb{R}^N\right\}, \tag{2.7}$$

where $c^A$ is the linear operator defined by $c^A = \sum_{n=0}^\infty \frac{(\ln c)^n A^n}{n!}$. Xiao (2009) proved that the Gaussian random field $X_0$ satisfies Conditions (C1) and (C2), and characterized many sample path properties of the corresponding $(N, d)$-Gaussian field $X$ in terms of $(H_1, \ldots, H_N)$ explicitly.

We remark that all centered stationary Gaussian random fields can also be treated using the above framework. In fact, if $Y = \{Y(t), t \in \mathbb{R}^N\}$ is a centered, real-valued stationary Gaussian random field, it can be represented as $Y(t) = \int_{\mathbb{R}^N} e^{i\langle t, \lambda\rangle}\, W(d\lambda)$. Thus the random field $X_0$ defined by

$$X_0(t) = Y(t) - Y(0) = \int_{\mathbb{R}^N} \left(e^{i\langle t,\lambda\rangle} - 1\right) W(d\lambda), \quad \forall\, t \in \mathbb{R}^N$$

is Gaussian with stationary increments and $X_0(0) = 0$. Note that the spectral measure $F$ of $X_0$ in the sense of (2.4) is the same as the spectral measure [in the ordinary sense] of the stationary random field $Y$.

The main purpose of this section is to prove a sufficient condition for a general Gaussian random field $X_0$ with stationary increments to satisfy Conditions (C1) and (C2). In particular, this condition implies that $X_0$ is strongly locally nondeterministic in metric $\rho$.

To this end we first introduce some notation and state several lemmas. For any $\lambda \in \mathbb{R}^N$ and $h > 0$, we denote by $C(\lambda, h)$ the cube with side-length $2h$ and center $\lambda$, i.e.,

$$C(\lambda, h) = \left\{x \in \mathbb{R}^N : |x_j - \lambda_j| \le h,\, j = 1, \cdots, N\right\}.$$

For any $g \in L^2(\mathbb{R}^N)$, let $\widehat{g}(\lambda) = \int_{\mathbb{R}^N} e^{i\langle \lambda, x\rangle} g(x) dx$ be the Fourier transform of $g$ and let $L^2(C(0,T))$ denote the subspace of $g \in L^2(\mathbb{R}^N)$ whose support is contained in $C(0, T)$. In the following, Lemma 2.1 is Proposition 4 of Pitt (1975). Lemma 2.2 is taken from Xiao (2007), which is an extension of a result of Pitt (1978, p.326).

**Lemma 2.1** *Let* $\widetilde{\Delta}(d\lambda)$ *be a positive measure on* $\mathbb{R}^N$*. If, for some constant* $h > 0$*,* $\widetilde{\Delta}(d\lambda)$ *satisfies*

$$0 < \liminf_{\|\lambda\|\to\infty} \widetilde{\Delta}(C(\lambda, h)) \le \limsup_{\|\lambda\|\to\infty} \widetilde{\Delta}(C(\lambda, h)) < \infty, \tag{2.8}$$

*then, for every* $T > 0$ *satisfying* $ThN < \log 2$*, there exist positive and finite constants* $c_{2,2}$ *and* $c_{2,3}$ *such that*

$$c_{2,2} \int_{\mathbb{R}^N} |\widehat{\psi}(\lambda)|^2 d\lambda \le \int_{\mathbb{R}^N} |\widehat{\psi}(\lambda)|^2 \widetilde{\Delta}(d\lambda) \le c_{2,3} \int_{\mathbb{R}^N} |\widehat{\psi}(\lambda)|^2 d\lambda \tag{2.9}$$

*for all* $\psi \in L^2(C(0,T))$*.*

**Lemma 2.2** *Let* $\Delta_1(d\lambda)$ *be a positive measure on* $\mathbb{R}^N$ *with density function* $\Delta_1(\lambda)$*. If there exist constants* $c_{2,4} > 0$ *and* $\eta > 0$ *such that*

$$\Delta_1(\lambda) \ge \frac{c_{2,4}}{\|\lambda\|^\eta} \qquad \text{for all } \lambda \in \mathbb{R}^N \text{ with } \|\lambda\| \text{ large.} \tag{2.10}$$

*Then for any constants* $T > 0$ *and* $c_{2,5}$*, there exists a positive and finite constant* $c_{2,6}$ *such that for all functions* $g$ *of the form*

$$g(\lambda) = \sum_{j=1}^{n} a_j \left( e^{i\langle s^j, \lambda \rangle} - 1 \right), \tag{2.11}$$

*where* $a_j \in \mathbb{R}$ *and* $s^j \in C(0,T)$*, we have*

$$|g(\lambda)| \le c_{2,6} \|\lambda\| \cdot \left( \int_{\mathbb{R}^N} |g(\xi)|^2 \ \Delta_1(\xi) d\xi \right)^{1/2}$$

*for all* $\lambda \in \mathbb{R}^N$ *with* $\|\lambda\| \le c_{2,5}$*.*

Lemma 2.3 below is an extension of Proposition 8.4 of Pitt (1978). It allows us to connect the property of strong local nondeterminism of a Gaussian random field with a general spectral measure to that of a Gaussian random field with an absolutely continuous spectral measure, which has been studied in Xiao (2007, 2009).

**Lemma 2.3** *Let* $\Delta_2(d\lambda)$ *be a positive measure on* $\mathbb{R}^N$ *and suppose that for some* $h > 0$*,*

$$0 < \liminf_{\|\lambda\|\to\infty} \rho(0,\lambda)^{Q+2} \Delta_2(C(\lambda, h)) \le \limsup_{\|\lambda\|\to\infty} \rho(0,\lambda)^{Q+2} \Delta_2(C(\lambda, h)) < \infty. \tag{2.12}$$

*Then for any constant* $T > 0$ *with* $ThN < \log 2$*, there exist positive and finite constants* $c_{2,7}$ *and* $c_{2,8}$ *such that*

$$c_{2,7} \int_{\mathbb{R}^N} \frac{|g(\lambda)|^2}{(\sum_{j=1}^N |\lambda_j|^{H_j})^{Q+2}} d\lambda \le \int_{\mathbb{R}^N} |g(\lambda)|^2 \Delta_2(d\lambda) \le c_{2,8} \int_{\mathbb{R}^N} \frac{|g(\lambda)|^2}{(\sum_{j=1}^N |\lambda_j|^{H_j})^{Q+2}} d\lambda \tag{2.13}$$

*for all* $g(\lambda)$ *of the form (2.11).*

**Proof.** First we claim that there is a positive constant $c \le 1$ such that

$$\begin{aligned} c \int_{\mathbb{R}^N} \frac{|g(\lambda)|^2}{\left(\sum_{j=1}^N |\lambda_j|^{H_j}\right)^{Q+2}}\, d\lambda &\le \int_{\mathbb{R}^N} \frac{|g(\lambda)|^2}{\left(1+\sum_{j=1}^N |\lambda_j|^{H_j}\right)^{Q+2}}\, d\lambda \\ &\le \int_{\mathbb{R}^N} \frac{|g(\lambda)|^2}{\left(\sum_{j=1}^N |\lambda_j|^{H_j}\right)^{Q+2}}\, d\lambda \end{aligned} \tag{2.14}$$

for all functions $g$ of the form (2.11).

Clearly only the first inequality in (2.14) needs a proof. For this purpose, we split the first integral in (2.14) over $\{\lambda : \|\lambda\| \le c_{2,5}\}$ and $\{\lambda : \|\lambda\| > c_{2,5}\}$ and apply Lemma 2.2 with

$$\Delta_1(d\lambda) = \frac{d\lambda}{\left(1+\sum_{j=1}^N |\lambda_j|^{H_j}\right)^{Q+2}}$$

[which satisfies (2.10)] to derive

$$\begin{aligned} &\int_{\{\|\lambda\| \le c_{2,5}\}} \frac{|g(\lambda)|^2}{\left(\sum_{j=1}^N |\lambda_j|^{H_j}\right)^{Q+2}}\, d\lambda \\ &\le c_{2,6}^2 \int_{\{\|\lambda\| \le c_{2,5}\}} \frac{\|\lambda\|^2}{\left(\sum_{j=1}^N |\lambda_j|^{H_j}\right)^{Q+2}}\, d\lambda \cdot \int_{\mathbb{R}^N} |g(\xi)|^2\, \Delta_1(d\xi) \\ &= c_{2,9} \int_{\mathbb{R}^N} |g(\xi)|^2\, \Delta_1(d\xi), \end{aligned}$$

because the first integral in the second line is convergent. It follows from the above that

$$\begin{aligned} \int_{\mathbb{R}^N} \frac{|g(\lambda)|^2}{\left(\sum_{j=1}^N |\lambda_j|^{H_j}\right)^{Q+2}}\, d\lambda &\le c_{2,9} \int_{\mathbb{R}^N} \frac{|g(\lambda)|^2}{\left(1+\sum_{j=1}^N |\lambda_j|^{H_j}\right)^{Q+2}}\, d\lambda \\ &\qquad + \int_{\{\lambda : \|\lambda\| > c_{2,5}\}} \frac{|g(\lambda)|^2}{\left(\sum_{j=1}^N |\lambda_j|^{H_j}\right)^{Q+2}}\, d\lambda \\ &\le c_{2,10} \int_{\mathbb{R}^N} \frac{|g(\lambda)|^2}{\left(1+\sum_{j=1}^N |\lambda_j|^{H_j}\right)^{Q+2}}\, d\lambda. \end{aligned}$$

This verifies the first inequality in (2.14).

Next we take a constant $s > 0$ such that $(T+s)hN < \log 2$ and denote $T_1 = T + s$. Let $\varphi \in L^2(C(0,s))$ be a function with the following property

$$c_{2,11} \le |\widehat{\varphi}(\lambda)|^2 \cdot (1+\rho(0,\lambda))^{Q+2} \le c_{2,12} \tag{2.15}$$

for all $\lambda \in \mathbb{R}^N$, where $c_{2,11}$ and $c_{2,12}$ are positive and finite constants. Such a function $\varphi$ can be constructed as follows. Observe that the function $\lambda \mapsto (1+\rho(0,\lambda))^{-(Q+2)/2}$ is in $L^2(\mathbb{R}^N)$.

Hence it is the Fourier transform of a function $\kappa \in L^2(\mathbb{R}^N)$. For the constant $s > 0$ chosen above we consider the function

$$P_s(t) = \prod_{j=1}^N \left(1 - \frac{|t_j|}{s}\right)^+ \qquad \text{for all } t \in \mathbb{R}^N,$$

where $a^+ := \max(a\,, 0)$ for all real numbers $a$. Then the support of $P_s$ is $C(0, s)$. Recall that the Fourier transform of $P_s$ is

$$\widehat{P}_s(\xi) := 2^N \prod_{j=1}^N \frac{1 - \cos(s\xi_j)}{s\xi_j^2} \qquad \text{for all } \xi \in \mathbb{R}^N.$$

Define $\varphi(t) = \kappa(t)P_s(t)$. Then $\varphi \in L^1(C(0, s)) \cap L^2(C(0, s))$ and its Fourier transform is given by

$$\begin{aligned}\widehat{\varphi}(\lambda) &= \widehat{\kappa} \star \widehat{P}_s(\lambda)\\ &= \int_{\mathbb{R}^N} \frac{2^N}{\big(1 + \rho(0, \lambda - \xi)\big)^{(Q+2)/2}} \prod_{j=1}^N \frac{1 - \cos(s\xi_j)}{s\xi_j^2}\, d\xi.\end{aligned}$$

It is clear that $\widehat{\varphi}(\lambda) > 0$ for all $\lambda \in \mathbb{R}^N$. Writing

$$\widehat{\varphi}(\lambda) \cdot (1 + \rho(0, \lambda))^{(Q+2)/2} = \int_{\mathbb{R}^N} \frac{2^N\, (1 + \rho(0, \lambda))^{(Q+2)/2}}{\big(1 + \rho(0, \lambda - \xi)\big)^{(Q+2)/2}} \prod_{j=1}^N \frac{1 - \cos(s\xi_j)}{s\xi_j^2}\, d\xi$$

and using the dominated convergence theorem, we see that

$$\lim_{\|\lambda\| \to \infty} \widehat{\varphi}(\lambda) \cdot (1 + \rho(0, \lambda))^{(Q+2)/2} = 2^N \int_{\mathbb{R}^N} \prod_{j=1}^N \frac{1 - \cos(s\xi_j)}{s\xi_j^2}\, d\xi.$$

Hence (2.15) follows.

Now we continue with the proof of (2.13). Let

$$\widehat{\psi}(\lambda) := g(\lambda)\widehat{\varphi}(\lambda) = \sum_{j=1}^n a_j \big(e^{i\langle s^j, \lambda\rangle} - 1\big)\widehat{\varphi}(\lambda),$$

where $s^j \in C(0, T)$ for $j = 1, \ldots, n$. Since $\varphi \in L^1(C(0, s)) \cap L^2(C(0, s))$, we use the Fourier inversion formula to verify that $\psi \in L^2(C(0, T_1))$. Moreover, by (2.14) and (2.15), there is a constant $c \geq 1$ such that

$$c^{-1} \int_{\mathbb{R}^N} |g(\lambda)\widehat{\varphi}(\lambda)|^2 d\lambda \leq \int_{\mathbb{R}^N} \frac{|g(\lambda)|^2}{\Big(\sum_{j=1}^N |\lambda_j|^{H_j}\Big)^{Q+2}} d\lambda \leq c \int_{\mathbb{R}^N} |g(\lambda)\widehat{\varphi}(\lambda)|^2 d\lambda \tag{2.16}$$

for all functions $g$ of the form (2.11).

Consider the new positive measure $\widetilde{\Delta}(d\lambda)$ on $\mathbb{R}^N$ defined by $\widetilde{\Delta}(d\lambda) = |\widehat{\varphi}(\lambda)|^{-2}\Delta_2(d\lambda)$. It follows from (2.12) and (2.15) that

$$\liminf_{\|\lambda\|\to\infty} \widetilde{\Delta}(C(\lambda,h)) \geq c \liminf_{\|\lambda\|\to\infty} \rho(0,\lambda)^{Q+2}\Delta_2(C(\lambda,h)) > 0$$

and

$$\limsup_{\|\lambda\|\to\infty} \widetilde{\Delta}(C(\lambda,h)) \leq c \limsup_{\|\lambda\|\to\infty} \rho(0,\lambda)^{Q+2}\Delta_2(C(\lambda,h)) < \infty$$

Hence the measure $\widetilde{\Delta}(d\lambda)$ satisfies (2.8). We apply Lemma 2.1 to derive that

$$\begin{aligned} c_{2,2} \int_{\mathbb{R}^N} |g(\lambda)\widehat{\varphi}(\lambda)|^2 d\lambda &\leq \int_{\mathbb{R}^N} |g(\lambda)\widehat{\varphi}(\lambda)|^2 \widetilde{\Delta}(d\lambda) \\ &= \int_{\mathbb{R}^N} |g(\lambda)|^2 \Delta_2(\lambda) \leq c_{2,3} \int_{\mathbb{R}^N} |g(\lambda)\widehat{\varphi}(\lambda)|^2 d\lambda. \end{aligned}$$

for all functions $g$ of the form (2.11) provided $s^j \in C(0,T)$ for $j = 1, \ldots, n$. This and (2.16) yield (2.13). □

We are ready to prove the main result of this section.

**Theorem 2.4** *Let* $\{X_0(t), t \in \mathbb{R}^N\}$ *be a real-valued centered Gaussian random field with stationary increments and* $X_0(0) = 0$. *If for some constant* $h > 0$ *the spectral measure* $F$ *of* $X_0$ *satisfies*

$$0 < \liminf_{\|\lambda\|\to\infty} \rho(0,\lambda)^{Q+2} F(C(\lambda,h)) \leq \limsup_{\|\lambda\|\to\infty} \rho(0,\lambda)^{Q+2} F(C(\lambda,h)) < \infty, \tag{2.17}$$

*then for any* $T > 0$ *such that* $ThN < \log 2$, $X_0$ *satisfies Conditions (C1) and (C2) on* $C(0,T)$.

**Proof.** First we verify $X_0$ satisfies Condition (C1). For any $s,t \in C(0,T)$, we apply the stochastic representation of $X_0$ and Lemma 2.3 to write

$$\begin{aligned} \mathbb{E}\big(|X_0(s) - X_0(t)|^2\big) &= \int_{\mathbb{R}^N} \left|e^{i\langle s,\lambda\rangle} - e^{i\langle t,\lambda\rangle}\right|^2 F(d\lambda) \\ &\asymp \int_{\mathbb{R}^N} \frac{\left|e^{i\langle s,\lambda\rangle} - e^{i\langle t,\lambda\rangle}\right|^2}{\left(\sum_{j=1}^N |\lambda_j|^{H_j}\right)^{Q+2}} \, d\lambda. \end{aligned} \tag{2.18}$$

Since it has been proved in Xiao (2009) that

$$\int_{\mathbb{R}^N} \frac{\left|e^{i\langle s,\lambda\rangle} - e^{i\langle t,\lambda\rangle}\right|^2}{\left(\sum_{j=1}^N |\lambda_j|^{H_j}\right)^{Q+2}} \, d\lambda \asymp \rho(s,t)^2, \qquad \forall s,t \in C(0,T),$$

we conclude that $X_0$ satisfies (C1) on $C(0,T)$.

Now we prove that $X_0$ satisfies Condition (C2) on $C(0,T)$. Denote $r = \min_{0\le j\le n} \rho(u, t^j)$. It is sufficient to prove that for all $a_j \in \mathbb{R}$ $(1 \le j \le n)$ we have

$$\mathbb{E}\left(\left|X_0(u) - \sum_{j=1}^{n} a_j X_0(t^j)\right|^2\right) \ge c_{2,10}\, r^2 \tag{2.19}$$

and $c_{2,10}$ is a positive constant which is independent of $n$, $a_j$ and the choice of $\{t^j\}$ and $u$. Again by using the stochastic representation of $X_0$, the left hand side of (2.19) can be written as

$$\begin{aligned}&\mathbb{E}\left(\left|X_0(u) - \sum_{j=1}^{n} a_j X_0(t^j)\right|^2\right)\\ &= \int_{\mathbb{R}^N} \left| e^{i\langle u,\lambda\rangle} - 1 - \sum_{j=1}^{n} a_j \left(e^{i\langle t^j,\lambda\rangle} - 1\right)\right|^2 F(d\lambda).\end{aligned}$$

Note that the function inside the integral is of the form (2.11). We apply Lemma 2.3 to get

$$\begin{aligned}&\int_{\mathbb{R}^N} \left| e^{i\langle u,\lambda\rangle} - 1 - \sum_{j=1}^{n} a_j \left(e^{i\langle t^j,\lambda\rangle} - 1\right)\right|^2 F(d\lambda)\\ &\ge c_{2,7} \int_{\mathbb{R}^N} \left| e^{i\langle u,\lambda\rangle} - 1 - \sum_{j=1}^{n} a_j \left(e^{i\langle t^j,\lambda\rangle} - 1\right)\right|^2 \frac{d\lambda}{\left(\sum_{j=1}^{N} |\lambda_j|^{H_j}\right)^{Q+2}}.\end{aligned}$$

However, it has been proved in Theorem 3.2 of Xiao (2009) that the last integral is bounded from below by $c_{2,11} r^2$, and $c_{2,11}$ is a positive constant which is independent of $n$, $a_j$ and the choice of $\{t^j\}$ and $u$. This proves (2.19) and Theorem 2.4. □

Theorem 2.4 can be applied directly to Gaussian random fields with stationary increments and with discrete spectral measure (or of mixed form $F = F_{ac} + F_{dis}$). It is useful for analyzing many space-time Gaussian random fields in the literature; see Xue and Xiao (2011) and the references therein for some examples. In the following we give an example of Gaussian random field with discrete spectral measure $F$.

Let $\{\xi_n, n \in \mathbb{Z}^N\}$ and $\{\eta_n, n \in \mathbb{Z}^N\}$ be two independent sequences of i. i. d. $N(0,1)$ random variables, where $\mathbb{Z}$ is the set of integers. Let $\{a_n, n \in \mathbb{Z}^N\}$ be a sequence of real numbers such that

$$\sum_{n\in\mathbb{Z}^N} a_n^2 < \infty.$$

Then

$$Y(t) = \sum_{n\in\mathbb{Z}^N} a_n \big(\xi_n \cos\langle n,t\rangle + \eta_n \sin\langle n,t\rangle\big), \qquad t \in \mathbb{R}^N$$

is a centered stationary Gaussian random field with covariance function

$$\mathbb{E}(Y(t)Y(s)) = \sum_{n \in \mathbb{Z}^N} a_n^2 \cos \langle n, t-s \rangle .$$

Hence the spectral measure $F$ of $Y$ is supported on $\mathbb{Z}^N$ with $F(\{n\}) = a_n^2$. If we choose $\{a_n\}$ such that as $\|n\| \to \infty$,

$$a_n^2 \asymp \frac{1}{\left(\sum_{j=1}^N n_j^{H_j}\right)^{Q+2}},$$

then for any fixed constant $h > 1$, $F$ satisfies (2.17). Consider the Gaussian random field $\{X_0(t), t \in \mathbb{R}^N\}$ defined by $X_0(t) = Y(t) - Y(0)$. Theorem 2.4 implies that, for any constant $T > 0$ with $ThN < \log 2$, $\{X_0(t), t \in \mathbb{R}^N\}$ satisfies Conditions (C1) and (C2) on $C(0,T)$.

Consequently, many sample path properties of $Y$ such as uniform and local moduli of continuity, Chung's law of the iterated logarithm, existence and joint continuity of the local times can be derived from the results in Xiao (2009), Luan and Xiao (2010), and Meerschaert *et al.* (2011).

# 3 Exact Hausdorff measure function for the range $X([0,1]^N)$

In this section, we determine the exact Hausdorff measure function for the range of an $(N,d)$-Gaussian random field $X = \{X(t), t \in \mathbb{R}^N\}$ defined in (1.1), where $X_1, \ldots, X_d$ are independent copies of a real-valued, centered Gaussian random field $X_0$ with stationary increments, which satisfies Conditions (C1) and (C2).

First we recall briefly the definition of Hausdorff measure, an upper density theorem due to Rogers and Taylor (1961) and two useful inequalities for large and small tails of the supremum of Gaussian processes. Then we extend a result of Talagrand (1995) to anisotropic Gaussian random fields, which is applied to derive an upper bound for the $\varphi_1$-Hausdorff measure of $X([0,1]^N)$. Finally we prove a law of the iterated logarithm for the sojourn time of $X$ and derive a lower bound for the $\varphi_1$-Hausdorff measure of $X([0,1]^N)$.

## 3.1 Hausdorff measure

Let $\Phi$ be the class of functions $\phi : (0,\delta) \to (0,1)$ which are right continuous, monotone increasing with $\phi(0_+) = 0$ and such that there exists a finite constant $c_{3,1} > 0$ for which

$$\frac{\phi(2s)}{\phi(s)} \le c_{3,1}, \quad \text{for } 0 < s < \frac{1}{2}\delta.$$

For $\phi \in \Phi$, the $\phi$-Hausdorff measure of $E \subseteq \mathbb{R}^d$ is defined by

$$\phi\text{-}m(E) = \lim_{\epsilon \to 0} \inf \left\{ \sum_i \phi(2r_i) : \, E \subseteq \bigcup_{i=1}^{\infty} B(x_i, r_i), r_i < \epsilon \right\},$$

where $B(x,r)$ denotes the Euclidean open ball of radius $r$ centered at $x$. It is known that $\phi$-$m$ is a metric outer measure and every Borel set in $\mathbb{R}^d$ is $\phi$-$m$ measurable. We say that a function

$\phi$ is an exact Hausdorff measure function for $E$ if $0 < \phi\text{-}m(E) < \infty$. The Hausdorff dimension of $E$ is defined by

$$\begin{aligned} \dim_{\mathrm{H}} E &= \inf\{\alpha > 0; s^{\alpha}\text{-}m(E) = 0\} \\ &= \sup\{\alpha > 0; s^{\alpha}\text{-}m(E) = \infty\}. \end{aligned}$$

We refer to Falconer (1990) for more properties of Hausdorff measure and Hausdorff dimension.

The following lemma can be easily derived from the results in Rogers and Taylor (1961), which gives a way to get a lower bound for $\phi\text{-}m(E)$. For any Borel measure $\mu$ on $\mathbb{R}^d$ and $\phi \in \Phi$, the upper $\phi$-density of $\mu$ at $x \in \mathbb{R}^d$ is defined by

$$\overline{D}_{\mu}^{\phi}(x) = \limsup_{r \to 0} \frac{\mu(B(x,r))}{\phi(2r)}.$$

**Lemma 3.1** *For a given $\phi \in \Phi$ there exists a positive constant $c_{3,2}$ such that for any Borel measure $\mu$ on $\mathbb{R}^d$ and every Borel set $E \subseteq \mathbb{R}^d$, we have*

$$\phi\text{-}m(E) \geq c_{3,2}\, \mu(E) \inf_{x \in E} \{\overline{D}_{\mu}^{\phi}(x)\}^{-1}.$$

Now we recall some basic facts about Gaussian processes. Consider a set $S$ and a centered Gaussian process $\{Y(t), t \in S\}$. We provide $S$ with the following canonical pseudo-metric

$$d(s,t) = \|Y(s) - Y(t)\|_2,$$

where $\|Y\|_2 = (\mathbb{E}(Y^2))^{1/2}$. Denote by $N_d(S, \epsilon)$ the smallest number of open $d$-balls of radius $\epsilon$ needed to cover $S$ and let $D = \sup\{d(s,t) : s, t \in S\}$ be the $d$-diameter of $S$.

The following lemma is well known. It is a consequence of the Gaussian isoperimetric inequality and Dudley's entropy bound [see Talagrand (1995)].

**Lemma 3.2** *There exists a positive constant $c_{3,3}$ such that for all $u > 0$, we have*

$$\mathbb{P}\left\{ \sup_{s,t \in S} |Y(s) - Y(t)| \geq c_{3,3} \left( u + \int_0^D \sqrt{\log N_d(S, \epsilon)}\, d\epsilon \right) \right\} \leq \exp\left( -\frac{u^2}{D^2} \right).$$

**Lemma 3.3** *Consider a function $\Psi$ such that $N_d(S, \epsilon) \leq \Psi(\epsilon)$ for all $\epsilon > 0$. Assume that for some constant $c_{3,4} \geq 1$ and all $\epsilon > 0$ we have*

$$\Psi(\epsilon)/c_{3,4} \leq \Psi(\frac{\epsilon}{2}) \leq c_{3,4} \Psi(\epsilon).$$

*Then*

$$\mathbb{P}\left\{ \sup_{s,t \in S} |Y(s) - Y(t)| \leq u \right\} \geq \exp\big( - c_{3,5} \Psi(u) \big),$$

*where $c_{3,5} > 0$ is a constant depending only on $c_{3,4}$.*

This was proved in Talagrand (1993). It gives a general lower bound for the small ball probability of Gaussian processes.

## 3.2 Some basic estimates

Let $X_0 = \{X_0(t),\, t \in \mathbb{R}^N\}$ be a centered Gaussian random field with stationary increments and satisfying Conditions (C1) and (C2). Without loss of generality, we assume that $H_1, \ldots, H_N$ are ordered as

$$0 < H_1 \le H_2 \le \cdots \le H_N < 1. \tag{3.1}$$

In order to solve some dependence problems that are a major obstacle, we consider for any given $0 < a < b < \infty$ the random field

$$X_0(a,b,t) = \int_{a < \rho(0,\lambda) \le b} \left(e^{i\langle t,\lambda\rangle} - 1\right) W(d\lambda), \qquad t \in \mathbb{R}^N.$$

An essential fact is that if $0 < a < b < a' < b' < \infty$, then the Gaussian random fields $\{X_0(a,b,t), t \in \mathbb{R}^N\}$ and $\{X_0(a',b',t), t \in \mathbb{R}^N\}$ are independent.

Let $X_1(a,b,t), \cdots, X_d(a,b,t)$ be independent copies of $X_0(a,b,t)$ and let

$$X(a,b,t) = \left(X_1(a,b,t), \cdots, X_d(a,b,t)\right), \qquad t \in \mathbb{R}^N.$$

Then we have the following lemma. For convenience, we write $I = [0,1]^N$.

**Lemma 3.4** *Given any* $0 < a < b$ *and* $0 < \epsilon < r$, *we have*

$$\mathbb{P}\left\{ \sup_{t \in I: \rho(0,t) \le r} \|X(a,b,t)\| \le \epsilon \right\} \ge \exp\left(-c\left(\frac{r}{\epsilon}\right)^Q\right), \tag{3.2}$$

*where* $0 < c < \infty$ *is an absolute constant.*

**Proof.** It is sufficient to prove (3.2) for $X_0(a,b,t)$. Let $S = \{t \in I : \rho(0,t) \le r\}$ and define a distance $d$ on $S$ by

$$d(s,t) = \big\|X_0(a,b,s) - X_0(a,b,t)\big\|_2.$$

Then (C1) implies $d(s,t) \le c_{1,1} \sum_{i=1}^N |s_i - t_i|^{H_i}$ for all $s,t \in I$, independent of the choices of $0 < a < b$. It follows that

$$N_d(S,\epsilon) \le c\left(\frac{r}{\epsilon}\right)^Q.$$

By Lemma 3.3 we have

$$\mathbb{P}\left\{ \sup_{t \in I: \rho(0,t) \le r} |X_0(a,b,t)| \le \epsilon \right\} \ge \exp\left(-c\left(\frac{r}{\epsilon}\right)^Q\right).$$

This proves Lemma 3.4. □

The following truncation inequalities are extensions of those in Loéve (1977, p.209) for $N = 1$ and (3.4) and (3.5) in Xiao (1996) for $N > 1$ and $\rho$ being replaced by the Euclidean metric.

**Lemma 3.5** *There exist positive finite constants* $c_{3,6}$ *and* $c_{3,7}$ *such that the following hold.*

(i) *For any* $a > 0$ *and any* $t \in \mathbb{R}^N$ *with* $\rho(0,t)a \le 1/N$ *we have*

$$\int_{\{\lambda:\rho(0,\lambda)\le a\}} \langle t,\lambda\rangle^2 F(d\lambda) \le c_{3,6} \int_{\mathbb{R}^N} (1-\cos\langle t,\lambda\rangle) F(d\lambda). \tag{3.3}$$

(ii) *For all* $a > 0$

$$\int_{\{\lambda:\rho(0,\lambda)>a\}} F(d\lambda) \le c_{3,7} a^{-2}. \tag{3.4}$$

**Proof.** Notice that when $\rho(0,\lambda) \le a$, the condition $\rho(0,t)a \le 1/N$ implies that $|\langle t,\lambda\rangle| < 1$. It follows that

$$1-\cos\langle t,\lambda\rangle \ge \frac{\langle t,\lambda\rangle^2}{2}\left(1-\frac{\langle t,\lambda\rangle^2}{12}\right) \ge \frac{11}{24}\langle t,\lambda\rangle^2 .$$

Then for any $t \in \mathbb{R}^N$ with $\rho(0,t)a \le 1/N$ we have

$$\begin{aligned}\int_{\mathbb{R}^N}(1-\cos\langle t,\lambda\rangle)F(d\lambda) &\ge \frac{11}{24}\int_{\{\lambda:|\langle t,\lambda\rangle|\le 1\}} \langle t,\lambda\rangle^2 F(d\lambda)\\ &\ge \frac{11}{24}\int_{\{\lambda:\rho(0,\lambda)\le a\}} \langle t,\lambda\rangle^2 F(d\lambda).\end{aligned}$$

That is

$$\int_{\{\lambda:\rho(0,\lambda)\le a\}} \langle t,\lambda\rangle^2 F(d\lambda) \le \frac{24}{11}\int_{\mathbb{R}^N}(1-\cos\langle t,\lambda\rangle)F(d\lambda).$$

To prove (3.4), we make the following two claims:

(a). For any $u > 0$, if $\lambda_i \ne 0$ for $i = 1,\ldots,N$, then

$$\frac{1}{2^N u^Q}\int_{\prod_{i=1}^N [-u^{\frac{1}{H_i}}, u^{\frac{1}{H_i}}]} \cos\langle t,\lambda\rangle\, dt = \prod_{i=1}^N \frac{\sin\left(u^{\frac{1}{H_i}}\lambda_i\right)}{u^{\frac{1}{H_i}}\lambda_i}.$$

(b). For any $u > 0$,

$$\int_{\{\lambda:\rho(0,\lambda)>\frac{1}{u}\}} F(d\lambda) \le \frac{c}{2^N u^Q}\int_{\prod_{i=1}^N [-u^{\frac{1}{H_i}}, u^{\frac{1}{H_i}}]} dt \int_{\mathbb{R}^N}(1-\cos\langle t,\lambda\rangle)F(d\lambda).$$

Claim (a) is obviously true when $N = 1$. Suppose it is true for $N = k$, then for $N = k + 1$, we have

$$\begin{aligned}
&\frac{1}{2^{k+1}u^{\frac{1}{H_1}+\cdots+\frac{1}{H_{k+1}}}} \int_{\prod_{i=1}^{k}[-u^{\frac{1}{H_i}},u^{\frac{1}{H_i}}]} dt_1 \cdots dt_k \int_{[-u^{\frac{1}{H_{k+1}}},u^{\frac{1}{H_{k+1}}}]} \cos(t_1\lambda_1 + \cdots + t_{k+1}\lambda_{k+1}) dt_{k+1} \\
&= \frac{1}{2^{k}u^{\frac{1}{H_1}+\cdots+\frac{1}{H_{k}}}} \int_{\prod_{i=1}^{k}[-u^{\frac{1}{H_i}},u^{\frac{1}{H_i}}]} dt_1 \cdots dt_k \\
&\qquad \times \frac{\sin(t_1\lambda_1 + \cdots + t_k\lambda_k + u^{\frac{1}{H_{k+1}}}\lambda_{k+1}) - \sin(t_1\lambda_1 + \cdots + t_k\lambda_k - u^{\frac{1}{H_{k+1}}}\lambda_{k+1})}{2u^{\frac{1}{H_{k+1}}}\lambda_{k+1}} \\
&= \frac{1}{2^{k}u^{\frac{1}{H_1}+\cdots+\frac{1}{H_{k}}}} \int_{\prod_{i=1}^{k}[-u^{\frac{1}{H_i}},u^{\frac{1}{H_i}}]} \cos(t_1\lambda_1 + \cdots + t_k\lambda_k) dt_1 \cdots dt_k \frac{\sin u^{\frac{1}{H_{k+1}}}\lambda_{k+1}}{u^{\frac{1}{H_{k+1}}}\lambda_{k+1}} \\
&= \frac{\sin u^{\frac{1}{H_1}}\lambda_1}{u^{\frac{1}{H_1}}\lambda_1} \cdots \frac{\sin u^{\frac{1}{H_{k+1}}}\lambda_{k+1}}{u^{\frac{1}{H_{k+1}}}\lambda_{k+1}}.
\end{aligned}$$

Hence claim (a) is true for all $N \geq 1$.

By Fubini's theorem and claim (a), we have

$$\begin{aligned}
&\frac{1}{2^N u^Q} \int_{\prod_{i=1}^{N}[-u^{\frac{1}{H_i}},u^{\frac{1}{H_i}}]} dt \int_{\mathbb{R}^N} (1 - \cos\langle t, \lambda\rangle) F(d\lambda) \\
&= \int_{\mathbb{R}^N} \left[ \frac{1}{2^N u^Q} \int_{\prod_{i=1}^{N}[-u^{\frac{1}{H_i}},u^{\frac{1}{H_i}}]} (1 - \cos\langle t, \lambda\rangle) dt \right] F(d\lambda) \\
&= \int_{\mathbb{R}^N} \left( 1 - \prod_{i=1}^{N} \frac{\sin u^{\frac{1}{H_i}}\lambda_i}{u^{\frac{1}{H_i}}\lambda_i} \right) F(d\lambda) \\
&\geq \int_{\mathbb{R}^N \setminus \{\lambda : |\lambda_i| \leq (Nu)^{-\frac{1}{H_i}}, \forall i\}} \left( 1 - \prod_{i=1}^{N} \frac{\sin u^{\frac{1}{H_i}}\lambda_i}{u^{\frac{1}{H_i}}\lambda_i} \right) F(d\lambda) \\
&\geq c \int_{\mathbb{R}^N \setminus \{\lambda : |\lambda_i| \leq (Nu)^{-\frac{1}{H_i}}, \forall i\}} F(d\lambda) \\
&\geq c \int_{\{\lambda : \rho(0,\lambda) > \frac{1}{u}\}} F(d\lambda).
\end{aligned}$$

Hence claim (b) is verified.

Now we turn to the proof of (3.4). With claim (b), (2.4) and Condition (C1) in hand, we have for $a > 0$,

$$\begin{aligned}
\int_{\{\lambda : \rho(0,\lambda) > a\}} F(d\lambda) &\leq \frac{ca^Q}{2^N} \int_{\prod_{i=1}^{N}[-a^{-\frac{1}{H_i}},a^{-\frac{1}{H_i}}]} dt \int_{\mathbb{R}^N} (1 - \cos\langle t, \lambda\rangle) F(d\lambda) \\
&\leq \frac{ca^Q}{2^N} \int_{\prod_{i=1}^{N}[-a^{-\frac{1}{H_i}},a^{-\frac{1}{H_i}}]} \sum_{i=1}^{N} |t_i|^{2H_i} dt \\
&\leq ca^{-2}.
\end{aligned}$$

This finishes the proof of Lemma 3.5. □

Lemma 3.6 gives estimates on the small ball probability of the $(N,d)$-Gaussian random field $X$ in (1.1).

**Lemma 3.6** *There exist constants* $c_{3,8}$ *and* $c_{3,9}$ *such that for all* $0<\epsilon<r$,

$$\exp\bigg(-c_{3,8}\Big(\frac{r}{\epsilon}\Big)^Q\bigg)\le \mathbb{P}\bigg\{\sup_{t\in I:\rho(0,t)\le r}\|X(t)\|\le\epsilon\bigg\}\le \exp\bigg(-c_{3,9}\Big(\frac{r}{\epsilon}\Big)^Q\bigg). \tag{3.5}$$

**Proof.** Let $S=\{t\in I:\rho(0,t)\le r\}$. It follows from (C1) that for all $\epsilon\in(0,r)$,

$$N_\rho(S,\epsilon)\le c\prod_{i=1}^N\Big(\frac{r}{\epsilon}\Big)^{\frac{1}{H_i}}=c\Big(\frac{r}{\epsilon}\Big)^Q:=\psi(\epsilon).$$

Clearly $\psi(\epsilon)$ satisfies the condition in Lemma 3.3. Hence the lower bound in (3.5) follows from Lemma 3.3.

The proof of the upper bound in (3.5) is based on Condition (C2) and a conditioning argument and is similar to the proof of Theorem 5.1 in Xiao (2009) [see also Monrad and Rootzén(1995)]. We include it for the sake of completeness. Let $T=\prod_{i=1}^N\big[0,(\frac{r}{N})^{\frac{1}{H_i}}\big]$. Then $T\subseteq S$. We divide $T$ into

$$\ell:=\prod_{i=1}^N\bigg(\lfloor(\frac{r}{N\epsilon})^{\frac{1}{H_i}}\rfloor+1\bigg)\ge\Big(\frac{r}{N\epsilon}\Big)^Q$$

sub-rectangles of side-lengths $\epsilon^{1/H_i}$ $(i=1,\ldots,N)$, where $\lfloor x\rfloor$ is the largest integer no more than $x$. And denote the lower-left vertices of these rectangles (in any order) by $t_k$ $(k=1,\ldots,\ell)$. Then

$$\mathbb{P}\bigg\{\sup_{t\in S}\|X(t)\|\le\epsilon\bigg\}\le\mathbb{P}\bigg\{\sup_{1\le k\le\ell}\|X(t_k)\|\le\epsilon\bigg\}. \tag{3.6}$$

It follows from Condition (C2) that for every $1\le k\le\ell$

$$\mathrm{Var}\,(X_0(t_k)|X_0(t_i):1\le i\le k-1)\ge c_{1,2}\,\epsilon^2.$$

By this and Anderson's inequality for Gaussian measures [see Anderson (1995)], we have the following upper bound for the conditional probabilities

$$\mathbb{P}\big\{\|X(t_k)\|\le\epsilon\big|X(t_i):1\le i\le k-1\big\}\le\Phi\Big(\frac{1}{\sqrt{c_{1,2}}}\Big)^d, \tag{3.7}$$

where $\Phi(x)$ is the distribution function of a standard normal random variable. It follows from (3.6) and (3.7) that

$$\mathbb{P}\bigg\{\sup_{t\in S}\|X(t)\|\le\epsilon\bigg\}\le\Phi\Big(\frac{1}{\sqrt{c_{1,2}}}\Big)^{\ell d}\le\exp\bigg(-c_{3,9}\Big(\frac{r}{\epsilon}\Big)^Q\bigg).$$

Thus we obtain the upper bound in (3.5). □

The main estimate is given in the following proposition.

**Proposition 3.7** *There exist positive constants* $\delta_1$ *and* $c_{3,10}$ *such that for any* $0 < r_0 \le \delta_1$*, we have*

$$\begin{aligned}\mathbb{P}\left\{\exists\, r \in [r_0^2, r_0],\ \sup_{t\in I:\rho(0,t)\le r}\|X(t)\| \le c_{3,10}\, r\left(\log\log\frac{1}{r}\right)^{-1/Q}\right\}& \\ \ge 1-\exp\left(-\left(\log\frac{1}{r_0}\right)^{1/2}\right).&\end{aligned}\tag{3.8}$$

**Proof.** Though the main idea of the proof is similar to the proof of Proposition 4.1 in Talagrand (1995), some modifications are needed to characterize the anisotropic nature of $X$. Let $U > 1$ be a number whose value will be determined later. For $k \ge 0$, let $r_k = r_0 U^{-2k}$. Consider the largest integer $k_0$ such that

$$k_0 \le \frac{\log(1/r_0)}{2\log U}.\tag{3.9}$$

Thus, for $k \le k_0$ we have $r_0^2 \le r_k \le r_0$. It thereby suffices to prove that

$$\mathbb{P}\left\{\exists k \le k_0,\ \sup_{t\in I:\rho(0,t)\le r_k}\|X(t)\| \le c\, r_k\left(\log\log\frac{1}{r_k}\right)^{-1/Q}\right\} \ge 1-\exp\left(-\left(\log\frac{1}{r_0}\right)^{1/2}\right).$$

Let $a_k = r_0^{-1}U^{2k-1}$ and we define for $k = 0, 1, \cdots$

$$X_{0,k}(t) = X_0(a_k, a_{k+1}, t)$$

and

$$\widehat{X}_k(t) = \big(X_{1,k}(t), \cdots, X_{d,k}(t)\big),$$

where $X_{1,k}(t), \cdots, X_{d,k}(t)$ are independent copies of $X_{0,k}(t)$. Furthermore, we assume $X_1 - X_{1,k}, \cdots, X_d - X_{d,k}$ are independent copies of $X_0 - X_{0,k}$. We note that the Gaussian random fields $\widehat{X}_0, \widehat{X}_1, \cdots$ are independent. By Lemma 3.4 we can find a constant $c_{3,11} > 0$ such that, if $r_0$ is small enough, then for each $k \ge 0$

$$\begin{aligned}&\mathbb{P}\left\{\sup_{t\in I:\rho(0,t)\le r_k}\|\widehat{X}_k(t)\| \le c_{3,11}\, r_k\left(\log\log\frac{1}{r_k}\right)^{-1/Q}\right\}\\ &\ge \exp\left(-\frac{1}{4}\log\log\frac{1}{r_k}\right) = \frac{1}{(\log 1/r_k)^{\frac14}}\\ &\ge \frac{1}{(2\log 1/r_0)^{\frac14}}.\end{aligned}\tag{3.10}$$

By independence,

$$\mathbb{P}\left\{\exists k \le k_0,\ \sup_{t\in I:\rho(0,t)\le r_k}\|\widehat{X}_k(t)\| \le c_{3,11} r_k\left(\log\log\frac{1}{r_k}\right)^{-1/Q}\right\}$$

$$
\begin{aligned}
&\geq 1-\left(1-\frac{1}{(2\log 1/r_0)^{1/4}}\right)^{k_0}\\
&\geq 1-\exp\left(-\frac{k_0}{(2\log 1/r_0)^{1/4}}\right), \qquad (3.11)
\end{aligned}
$$

where the last inequality follows from the elementary inequality $1-x\leq e^{-x}$ for all $x\geq 0$.

Let $\beta=\min\{\frac{1}{H_N}-1,2\}$. We claim that for any $u\geq cr_kU^{-\frac{\beta}{2}}\sqrt{\log U}$,

$$
\mathbb{P}\left\{\sup_{t\in I:\rho(0,t)\leq r_k}\|X(t)-\widehat{X}_k(t)\|\geq u\right\}\leq \exp\left(-\frac{u^2}{cr_k^2U^{-\beta}}\right). \qquad (3.12)
$$

To see this, it's enough to prove that (3.12) holds for $X_0-X_{0,k}$. Consider $S=\{t\in I:\rho(0,t)\leq r_k\}$ and on $S$ the distance

$$
d(s,t)=\big\|(X_0(s)-X_{0,k}(s))-(X_0(t)-X_{0,k}(t))\big\|_2.
$$

Then $d(s,t)\leq c\sum_{i=1}^N|s_i-t_i|^{H_i}$ and $N_d(S,\epsilon)\leq c(\frac{r_k}{\epsilon})^Q$. Now we estimate the diameter $D$ of $S$. For any $t\in S$,

$$
\begin{aligned}
&\mathbb{E}\left[(X_0(t)-X_{0,k}(t))^2\right]=2\int_{\{\lambda:\rho(0,\lambda)\leq a_k\}\cup\{\lambda:\rho(0,\lambda)>a_{k+1}\}}(1-\cos\langle t,\lambda\rangle)F(d\lambda)\\
&\leq 2\int_{\{\lambda:\rho(0,\lambda)\leq a_k\}}(1-\cos\langle t,\lambda\rangle)F(d\lambda)+4\int_{\{\lambda:\rho(0,\lambda)>a_{k+1}\}}F(d\lambda)\\
&=: I_1+I_2. \qquad (3.13)
\end{aligned}
$$

The second term is easy to estimate: By Lemma 3.5,

$$
I_2\leq ca_{k+1}^{-2}. \qquad (3.14)
$$

For the first term $I_1$, we use the elementary inequality $1-\cos\langle t,\lambda\rangle\leq\frac{1}{2}\langle t,\lambda\rangle^2$ to derive that for all $t\in S$

$$
\begin{aligned}
I_1 &\leq \int_{\{\lambda:\rho(0,\lambda)\leq a_k\}}\langle t,\lambda\rangle^2F(d\lambda)\\
&= N^{\frac{2}{H_1}}U^{-\frac{1}{H_N}}\int_{\{\lambda:\rho(0,\lambda)\leq a_k\}}\Big\langle\frac{U^{\frac{1}{2H_N}}}{N^{\frac{1}{H_1}}}t,\lambda\Big\rangle^2F(d\lambda)\\
&= cU^{-\frac{1}{H_N}}\int_{\{\lambda:\rho(0,\lambda)\leq a_k\}}\langle t',\lambda\rangle^2F(d\lambda),
\end{aligned}
$$

where $t'=U^{\frac{1}{2H_N}}N^{-\frac{1}{H_1}}t$. Since $\rho(0,t')\leq\frac{1}{N}U^{\frac{1}{2}}\rho(0,t)\leq\frac{1}{N}U^{\frac{1}{2}}r_k<\frac{1}{Na_k}$, it follows from Lemma 3.5 and (C1) that

$$
I_1\leq cU^{-\frac{1}{H_N}}\rho(0,t')^2\leq cU^{1-\frac{1}{H_N}}\rho(0,t)^2\leq cr_k^2U^{-(\frac{1}{H_N}-1)}. \qquad (3.15)
$$

With (3.13), (3.14) and (3.15) in hand, the diameter of $S$ satisfies

$$
D^2 \leq c\left[r_k^2U^{-(\frac{1}{H_N}-1)}+a_{k+1}^{-2}\right]
$$

$$
\begin{aligned}
&\leq \; cr_k^2\left[U^{-(\frac{1}{H_N}-1)}+U^{-2}\right]\\
&\leq \; cr_k^2U^{-\beta}, \qquad (3.16)
\end{aligned}
$$

where $\beta=\min\{\frac{1}{H_N}-1,2\}$. Some simple calculations yield

$$
\begin{aligned}
\int_0^D \sqrt{\log N_d(S,\epsilon)}d\epsilon \;&\leq\; c\int_0^{cr_kU^{-\frac{\beta}{2}}}\sqrt{\log\frac{r_k}{\epsilon}}\,d\epsilon\\
&\leq\; cr_kU^{-\frac{\beta}{2}}\sqrt{\log U}. \qquad (3.17)
\end{aligned}
$$

Hence we use Lemma 3.2 and (3.17) to derive that for any $u\geq cr_kU^{-\frac{\beta}{2}}\sqrt{\log U}$,

$$
\mathbb{P}\left\{\sup_{\rho(0,t)\leq r_k}|X_0(t)-X_{0,k}(t)|\geq u\right\}\leq\exp\left(-\frac{u^2}{cr_k^2U^{-\beta}}\right). \qquad (3.18)
$$

Thus we have proved (3.12).

Now we continue our proof of (3.8). Let $U=(\log 1/r_0)^{1/\beta}$. We see that for $r_0>0$ small

$$
U^{\beta/2}\left(\log U\right)^{-1/2}\geq\left(\log\log\frac{1}{r_0}\right)^{1/Q}.
$$

Take $u=c_{3,11}r_k(\log\log 1/r_0)^{-1/Q}$. It follows from (3.12) that

$$
\begin{aligned}
&\mathbb{P}\left\{\sup_{t\in I:\rho(0,t)\leq r_k}\|X(t)-\widehat{X}_k(t)\|\geq c_{3,11}r_k\left(\log\log\frac{1}{r_0}\right)^{-1/Q}\right\}\\
&\qquad\leq\exp\left(-\frac{U^\beta}{c_{3,12}\left(\log\log 1/r_0\right)^{2/Q}}\right).
\end{aligned}
$$

Combining this with (3.11), we get

$$
\begin{aligned}
&\mathbb{P}\left\{\exists k\leq k_0,\ \sup_{\rho(0,t)\leq r_k}\|X(t)\|\leq 2c_{3,11}r_k\left(\log\log\frac{1}{r_k}\right)^{-1/Q}\right\}\\
&\qquad\geq 1-\exp\left(-\frac{k_0}{(2\log 1/r_0)^{1/4}}\right)\\
&\qquad\qquad-k_0\exp\left(-\frac{U^\beta}{c_{3,12}\left(\log\log 1/r_0\right)^{2/Q}}\right). \qquad (3.19)
\end{aligned}
$$

We recall that

$$
\frac{\log\left(1/r_0\right)}{4\log U}\leq k_0\leq\log\frac{1}{r_0}.
$$

Then the right-hand side of (3.19) is at least $1-\exp(-(\log 1/r_0)^{1/2})$ when $r_0>0$ is small enough. This completes the proof. □

### 3.3 Upper bound for the Hausdorff measure of the range

We start with the following result on the uniform modulus of continuity of $X_0$. See, e.g., Xiao (2009). More precise result can be found in Meerschaert *et al.* (2011).

**Lemma 3.8** *Let $X_0 = \{X_0(t), t \in \mathbb{R}^N\}$ be a centered Gaussian random field with values in $\mathbb{R}$. If Condition (C1) is satisfied, then there exists a positive and finite constant $c_{3,13}$ such that*

$$\limsup_{\|\varepsilon\|\to 0} \frac{\sup_{t\in[0,1]^N,\, s\in[0,\varepsilon]} |X_0(t+s) - X_0(t)|}{\rho(0,\varepsilon)\sqrt{\log(1+\rho(0,\varepsilon)^{-1})}} \le c_{3,13}, \quad a.s. \tag{3.20}$$

Now we derive an upper bound for the Hausdorff measure of $X([0,1]^N)$.

**Theorem 3.9** *If $d > Q$, then there exists a constant $c_{3,14} > 0$ such that*

$$\varphi_1\text{-}m(X([0,1]^N)) \le c_{3,14} \qquad a.s., \tag{3.21}$$

*where $\varphi_1(r) = r^Q \log\log 1/r$.*

**Proof.** For $k \ge 1$, consider the set

$$\begin{aligned} R_k \ = \ \Big\{ & t \in [0,1]^N : \exists\, r \in [2^{-2k}, 2^{-k}] \text{ such that} \\ & \sup_{s\in I:\rho(s,t)\le r} \|X(s) - X(t)\| \le c_{3,10}\, r(\log\log \frac{1}{r})^{-1/Q} \Big\}. \end{aligned} \tag{3.22}$$

By Proposition 3.7 we have

$$\mathbb{P}\{t \in R_k\} \ge 1 - \exp(-\sqrt{k/2}).$$

Denote by $L_N$ the Lebesgue measure on $\mathbb{R}^N$. It follows from Fubini's theorem that $\mathbb{P}(\Omega_0) = 1$, where

$$\Omega_0 = \Big\{\omega : \ L_N(R_k) \ge 1 - \exp(-\sqrt{k}/4) \text{ infinitely often}\Big\}.$$

On the other hand, by Lemma 3.8, there exists an event $\Omega_1$ such that $\mathbb{P}(\Omega_1) = 1$ and for all $\omega \in \Omega_1$, there exists $n_1 = n_1(\omega)$ large enough such that for all $n \ge n_1$ and any rectangle $I_n$ of side-lengths $2^{-n/H_i} (i = 1, \cdots, N)$ that meets $[0,1]^N$, we have

$$\sup_{s,t\in I_n} \|X(t) - X(s)\| \le c2^{-n}\sqrt{\log[1 + (N2^{-n})^{-1}]} \le c2^{-n}\sqrt{n}. \tag{3.23}$$

Now for a fixed $\omega \in \Omega_0 \cap \Omega_1$, we show that $\varphi_1\text{-}m(X([0,1]^N)) \le c_{3,14} < \infty$. Consider $k \ge 1$ such that

$$L_N(R_k) \ge 1 - \exp(-\sqrt{k}/4).$$

For any $n \ge 1$, we divide $[0,1]^N$ into $2^{nQ}$ disjoint (half-open and half closed) rectangles of side-lengths $2^{-n/H_i} (i = 1, \cdots, N)$. Denote by $I_n(x)$ the rectangle of side-lengths $2^{-n/H_i} (i =$

$1, \cdots, N)$ containing $x$. For any $x \in R_k$ we can find the smallest integer $n$ with $k \le n \le 2k + \ell_0$ (where $\ell_0$ depends on $N$ only) such that

$$\sup_{s,t \in I_n(x)} \|X(t) - X(s)\| \le c2^{-n}(\log\log 2^n)^{-1/Q}. \tag{3.24}$$

Thus we have

$$R_k \subseteq V = \bigcup_{n=k}^{2k+\ell_0} V_n$$

and each $V_n$ is a union of rectangles $I_n(x)$ satisfying (3.24). Clearly $X(I_n(x))$ can be covered by a ball of radius

$$\rho_n = c2^{-n}(\log\log 2^n)^{-1/Q}.$$

Since $\varphi_1(2\rho_n) \le c2^{-nQ} = cL_N(I_n)$, we obtain

$$\sum_{n=k}^{k+\ell_0} \sum_{I_n \in V_n} \varphi_1(2\rho_n) \le \sum_n \sum_{I_n \in V_n} cL_N(I_n) = cL_N(V) \le c. \tag{3.25}$$

Thus $X(V)$ is contained in the union of a family of balls $B_n$ of radius $\rho_n$ with $\sum_n \varphi_1(2\rho_n) \le c$.

On the other hand, $[0,1]^N \backslash V$ is contained in a union of rectangles of side-lengths $2^{-q/H_i} (i = 1, \cdots, N)$ where $q = 2k + \ell_0$, none of which meets $R_k$. There can be at most

$$2^{Qq} L_N([0,1]^N \backslash V) \le c2^{Qq} \exp(-\sqrt{k}/4)$$

such rectangles. Since $\omega \in \Omega_1$, (3.23) implies that, for each of these rectangles $I_q$, $X(I_q)$ is contained in a ball of radius $c2^{-q}\sqrt{q}$. Thus $X([0,1]^N \backslash V)$ can be covered by a family $B_n$ of balls of radius $\rho_n = c2^{-q}\sqrt{q}$ such that

$$\sum_n \varphi_1(2\rho_n) \le (c2^{Qq} \exp(-\sqrt{k}/4)) \cdot (c2^{-qQ} q^{Q/2} \log\log(c2^q q^{-1/2})) \le 1 \tag{3.26}$$

for $k$ large enough. Since $k$ can be arbitrarily large, Theorem 3.9 follows from (3.25) and (3.26). □

## 3.4 Lower bound for the Hausdorff measure of the range

**Theorem 3.10** *If $d > Q$, then there exists a constant $c_{3,15} > 0$ such that*

$$\varphi_1\text{-}m(X([0,1]^N)) \ge c_{3,15} \qquad a.s., \tag{3.27}$$

*where $\varphi_1(r) = r^Q \log\log 1/r$.*

In order to prove Theorem 3.10, we first study the asymptotic behavior of the sojourn measure of $X$. For any $r > 0$ and $y \in \mathbb{R}^d$, define

$$T_y(r) = \int_I \mathbf{1}_{\{\|X(t)-y\| \le r\}}\, dt,$$

the sojourn time of $X$ in the ball $B(y, r)$. If $y = 0$, we write $T(r)$ for $T_0(r)$.

**Lemma 3.11** *If $d > Q$, then there is a finite constant $c_{3,16}$ such that*

$$\mathbb{E}\left(T(r)^n\right) \le c_{3,16}^n n!\, r^{Qn} \tag{3.28}$$

*for all for all integers $n \ge 1$ and $0 < r < 1$.*

**Proof.** For $n = 1$, by Fubini's theorem and (C1) we have

$$\begin{aligned}
\mathbb{E}\big(T(r)\big) &= \int_I \mathbb{P}\left\{\|X(t)\| < r\right\} dt \\
&\le \int_I \min\left\{1, c\Big(\frac{r}{\rho(0,t)}\Big)^d\right\} dt \\
&= \int_{\{t\in I:\rho(0,t)\le cr\}} dt + c\int_{\{t\in I:\rho(0,t)> cr\}} \left(\frac{r}{\rho(0,t)}\right)^d dt \\
&=: J_1 + J_2.
\end{aligned}$$

The first term is easy to estimate:

$$J_1 \le c\prod_{i=1}^N r^{\frac{1}{H_i}} = cr^Q. \tag{3.29}$$

For the second term, we use the following elementary fact: Given positive constants $\beta$ and $\gamma$, there exists a finite constant $c_{3,17}$ such that for all $a > 0$,

$$\int_0^\infty \frac{dx}{(a+x^\beta)^\gamma} = \begin{cases} c_{3,17}\, a^{-(\gamma-\frac{1}{\beta})} & \text{if } \ \beta\gamma > 1, \\ +\infty & \text{if } \ \beta\gamma \le 1. \end{cases} \tag{3.30}$$

Since $\rho(0,t) > cr$ implies that $t_{j_0} \ge cr^{1/H_{j_0}}$ for some $j_0 \in \{1, \cdots, N\}$, without loss of generality we assume $j_0 = 1$. Then using (3.30) $(N-1)$ times, we obtain

$$\begin{aligned}
J_2 &\le cr^d \int_{cr^{\frac{1}{H_1}}}^1 dt_1 \int_{[0,1]^{N-1}} \frac{dt_2, \cdots, dt_N}{(\sum_{i=1}^N t_i^{H_i})^d} \\
&\le cr^d \int_{cr^{\frac{1}{H_1}}}^1 dt_1 \int_{[0,1]^{N-2}} \frac{dt_2, \cdots, dt_{N-1}}{(\sum_{i=1}^{N-1} t_i^{H_i})^{d-\frac{1}{H_N}}} \\
&\le cr^d \int_{cr^{\frac{1}{H_1}}}^1 \frac{dt_1}{(t_1^{H_1})^{d-\sum_{i=2}^N \frac{1}{H_i}}} \\
&\le c\, r^Q,
\end{aligned} \tag{3.31}$$

where the last step follows from the assumption that $d > Q$. It follows from (3.29) and (3.31) that

$$\mathbb{E}\big(T(r)\big) \le cr^Q. \tag{3.32}$$

For $n \ge 2$,

$$\mathbb{E}(T(r)^n) = \int_{I^n} \mathbb{P}\{\|X(t^j)\| < r, 1 \le j \le n\} dt^1 \cdots dt^n. \tag{3.33}$$

Consider $t^1, \cdots, t^n \in I$ satisfying

$$t^j \neq 0, \quad \text{for } j = 1, \cdots, n \ \text{ and } \ t^j \neq t^k \ \text{ for } \ j \neq k.$$

By Condition (C2), we have

$$\mathrm{Var}\big(X_0(t^n)\big|X_0(t^1), \cdots, X_0(t^{n-1})\big) \geq c_{1,2} \min_{0 \leq k \leq n-1} \rho(t^n, t^k)^2, \tag{3.34}$$

where $t^0 = 0$. Since conditional distributions in Gaussian processes are still Gaussian, (3.34) and Anderson's inequality yield that for all $x^1, \ldots, x^{n-1} \in \mathbb{R}^d$,

$$\begin{aligned} &\mathbb{P}\Big\{\|X(t^n)\| < r\big|X(t^1) = x^1, \cdots, X(t^{n-1}) = x^{n-1}\Big\} \\ &\qquad \leq c \min\bigg\{1, \ \bigg(\frac{r}{\min\limits_{0 \leq k \leq n-1} \rho(t^n, t^k)}\bigg)^d\bigg\}. \end{aligned} \tag{3.35}$$

It follows from (3.35) and an argument similar to the proof of (3.32) that

$$\begin{aligned} &\int_I \mathbb{P}\Big\{\|X(t^n)\| < r\big|X(t^1) = x^1, \cdots, X(t^{n-1}) = x^{n-1}\Big\} dt^n \\ &\qquad \leq c \int_I \sum_{k=0}^{n-1} \min\bigg\{1, c\Big(\frac{r}{\rho(t^n, t^k)}\Big)^d\bigg\}\, dt^n \\ &\qquad \leq c\, n \int_I \min\bigg\{1, c\Big(\frac{r}{\rho(0, t^n)}\Big)^d\bigg\}\, dt^n \\ &\qquad \leq c\, n r^Q. \end{aligned} \tag{3.36}$$

Combining (3.33) and (3.36), we obtain

$$\begin{aligned} \mathbb{E}\big(T(r)^n\big) &\leq cnr^Q \int_{I^{n-1}} \mathbb{P}\left\{\|X(t^1)\| < r, \cdots, \|X(t^{n-1})\| < r\right\} dt^1 \cdots dt^{n-1} \\ &= cnr^Q \mathbb{E}\big(T(r)^{n-1}\big). \end{aligned}$$

Hence the inequality (3.28) follows from this and induction. □

Let $0 < b < 1/c_{3,16}$. Then by (3.28) we have

$$\mathbb{E}\left(\exp\left(br^{-Q}T(r)\right)\right) \leq \sum_{n=0}^{\infty} (c_{3,16}\, b)^n < \infty. \tag{3.37}$$

This and the exponential Chebychev's inequality imply that for any constant $0 < b < 1/c_{3,16}$,

$$\mathbb{P}\big\{T(r) \geq r^Q u\big\} \leq \frac{e^{-bu}}{1 - c_{3,16}\, b} \tag{3.38}$$

for all $u > 0$.

The following is a law of the iterated logarithm for the sojourn measure of $X$.

**Proposition 3.12** *For every $\tau \in I$, we have*

$$\limsup_{r\to 0} \frac{T_{X(\tau)}(r)}{\varphi_1(r)} \le c_{3,16}, \quad a.s. \tag{3.39}$$

**Proof.** Since $\{X(t), t \in \mathbb{R}^N\}$ has stationary increments, it is sufficient to consider $\tau = 0$. Then (3.39) follows from (3.38), the Borel-Cantelli lemma and a monotonicity argument in a standard way. □

**Proof of Theorem 3.10.** We can prove this theorem by using Lemma 3.1 and Proposition 3.12, in the same way as that of Theorem 4.1 in Xiao (1996). □

**Proof of Theorem 1.1.** It follows immediately from Theorems 3.9 and 3.10. □

# References

[1] Adler, R. J. (1981), *The Geometry of Random Fields.* Wiley, New York.

[2] Anderson, T. W. (1955), The integral of a symmetric unimodal function over a symmetric convex set and some probability inequalities. *Proc. Amer. Math. Soc.* **6**, 170-176.

[3] Ayache, A. and Xiao, Y. (2005), Asymptotic growth properties and Hausdorff dimension of fractional Brownian sheets. *J. Fourier Anal. Appl.* **11**, 407–439.

[4] Baraka, D. and Mountford, T. (2008), A law of iterated logarithm for fractional Brownian motions. *Séminaire de Probabilités XLI. Lecture Notes in Math.* **1934**, 161–179, Springer, Berlin.

[5] Baraka, D. and Mountford, T. (2011), The exact Hausdorff measure of the zero set of fractional Brownian motion. *J. Theor. Probab.* **24**, 271-293.

[6] Benassi, A., Jaffard, S. and Roux, D. (1997), Elliptic Gaussian random Processes. *Rev. Mat. Iberoamericana* **13**, 19–90.

[7] Berg, C. and Forst, G. (1975), *Potential Theory on Locally Compact Abelian Groups.* Sringer-Verlag, New York-Heidelberg.

[8] Berman, S. M. (1973), Local nondeterminism and local times of Gaussian processes. *Indiana Univ. Math. J.* **23**, 69–94.

[9] Berman, S. M. (1988), Spectral conditions for local nondeterminism. *Stochastic Process. Appl.* **27**, 73–84.

[10] Biermé, H., Lacaux, C. and Xiao, Y. (2009), Hitting probabilities and the Hausdorff dimension of the inverse images of anisotropic Gaussian random fields. *Bull. London Math. Soc.* **41**, 253–273.

[11] Falconer, K. J. (1990), *Fractal Geometry – Mathematical Foundations and Applications.* Wiley & Sons, New York.

[12] Istas, J. (2005), Spherical and hyperbolic fractional Brownian motion. *Elec. Comm. Probab.* **10**, 254–262.

[13] Kahane, J.-P. (1985), *Some Random Series of Functions.* 2nd edition, Cambridge University Press, Cambridge.

[14] Loéve, L. (1977), *Probability Theory I.* Springer-Verlag, New York.

[15] Luan, N. and Xiao, Y. (2010), Chung's law of the iterated logarithm for anisotropic Gaussian random fields. *Statist. Probab. Lett.* **80**, 1886–1895.

[16] Meerschaert, M. M., Wang, W. and Xiao, Y. (2011), Fernique-type inequalities and moduli of continuity of anisotropic Gaussian random fields. *Trans. Amer. Math. Soc.*, to appear.

[17] Monrad, D. and Rootzén, H. (1995), Small values of Gaussian processes and functional laws of the iterated logarithm. *Proba. Theory Relat. Fields* **101**, 173–192.

[18] Nualart, E. and Viens, F. (2009), The fractional stochastic heat equation on the circle: time regularity and potential theory. *Stoch. Process. Appl.* **119**, 1505–1540.

[19] Pitt, L. D. (1975), Stationary Gaussian Markov fields on $\mathbb{R}^d$ with a deterministic component. *J. Multivar. Anal.* **5**, 300–311.

[20] Pitt, L. D. (1978), Local times for Gaussian vector fields. *Indiana Univ. Math. J.* **27**, 309–330.

[21] Rogers, C. A. and Taylor, S. J. (1961), Functions continuous and singular with respect to a Hausdorff measure. *Mathematika* **8**, 1–31.

[22] Talagrand, M. (1993), New Gaussian estimates for enlarged balls. *Geom. Funct. Anal.* **3**, 502–526.

[23] Talagrand, M. (1995), Hausdorff measure of trajectories of multiparameter fractional Brownian motion. *Ann. Probab.* **23**, 767–775.

[24] Talagrand, M. (1998), Multiple points of trajectories of multiparameter fractional Brownian motion. *Probab. Theory Relat. Fields* **112**, 545–563.

[25] Tindel, S., Tudor, C. A. and Viens, F. (2004), Sharp Gaussian regularity on the circle, and applications to the fractional stochastic heat equation. *J. Funct. Anal.* **217**, 280–313.

[26] Wu, D. and Xiao, Y. (2009), Uniform Hausdorff dimension results for Gaussian random fields. *Sci. in China, Ser. A* **52**, 1478–1496.

[27] Wu, D. and Xiao, Y. (2011), On local times of anisotropic Gaussian random fields. *Comm. Stoch. Anal.* **5**, 15–39.

[28] Xiao, Y. (1996), Hausdorff measure of the sample paths of Gaussian random fields. *Osaka J. Math.* **33**, 895–913.

[29] Xiao, Y. (1997a), Hausdorff dimension of the graph of fractional Brownian motion. *Math. Proc. Camb. Philos. Soc.* **122**, 565–576.

[30] Xiao, Y. (1997b), Hölder conditions for the local times and the Hausdorff measure of the level sets of Gaussian random fields. *Probab. Theory Relat. Fields* **109**, 129–157.

[31] Xiao, Y. (2007), Strong local nondeterminism of Gaussian random fields and its applications. In: *Asymptotic Theory in Probability and Statistics with Applications*, (T.-L. Lai, Q.-M. Shao and L. Qian, editors), pp. 136–176, Higher Education Press, Beijing.

[32] Xiao, Y. (2009), Sample path properties of anisotropic Gaussian random fields. In: *A Minicourse on Stochastic Partial Differential Equations,* (D. Khoshnevisan and F. Rassoul-Agha, editors), Lecture Notes in Math. **1962**, pp. 145–212, Springer, New York.

[33] Xue, Y. and Xiao, Y. (2011), Fractal and smoothness properties of anisotropic Gaussian models. *Frontiers Math. China*, to appear.

[34] Yaglom, A. M. (1957), Some classes of random fields in $n$-dimensional space, related to stationary random processes. *Th. Probab. Appl.* **2**, 273–320.

Nana Luan. School of Insurance and Economics, University of International Business and Economics, Beijing 100029, China
E-mail: `luannana318@gmail.com`

Yimin Xiao. Department of Statistics and Probability, A-413 Wells Hall, Michigan State University, East Lansing, MI 48824, U.S.A.
E-mail: `xiao@stt.msu.edu`
URL: `http://www.stt.msu.edu/~xiaoyimi`